


Exploring the (non-)coding genome through long-read sequencing

Thomas DERRIEN
(thomas.derrien@univ-rennes.fr)

CGT : Canine Genetics Team



IGDRion sequencing PF




The cell as an RNA machine




The cell as an RNA machine



The cell as an RNA machine




The cell as an RNA machine



Back2Basics

Gene


Pre-mRNA



Back2Basics

Gene

Pre-mRNA



Splicing

mRNA1


mRNA2

mRNA
(2 transcript isoforms)



mRNAs versus lncRNAs

Gene



lncRNA

(Transcript/Isoform)



lncRNA1




lncRNA2

lncRNAs are "mRNA-like" ...

- ✓ Transcribed by RNA Polymerase II
- ✓ Undergo splicing
- ✓ 5' capped
- ✓ Most have a polyA tail

mRNAs versus lncRNAs

Gene



lncRNA

(Transcript/Isoform)



lncRNA1



lncRNA2

lncRNAs are "mRNA-like" ...


- ✓ Transcribed by RNA Polymerase II
- ✓ Undergo splicing
- ✓ 5' capped
- ✓ Most have a polyA tail

... but are not mRNAs

⊘ No coding potential

mRNAs versus lncRNAs

Gene



lncRNA

(Transcript/Isoform)



lncRNA1



lncRNA2

lncRNAs are "mRNA-like" ...


- ✓ Transcribed by RNA Polymerase II
- ✓ Undergo splicing
- ✓ 5' capped
- ✓ Most have a polyA tail

... but are not mRNAs

- ✗ No coding potential
- ✗ Lowly expressed, highly tissue-specific
- ✗ Less conserved across species
- ✗ Often overlap Transposable Elements (TEs)

mRNAs versus lncRNAs

Gene



lncRNA

(Transcript/Isoform)



lncRNA1



lncRNA2

lncRNAs are "mRNA-like" ...

... but are not mRNAs



● Transcript >200nt with low coding potential



● Classified vs mRNA locations : lincRNA vs antisense



5' capped



Most have a polyA tail



Less conserved across species



Often overlap Transposable Elements (TEs)

lncRNA functions

lincRNAs act in the circuitry controlling pluripotency and differentiation

doi:10.1016/j.cell.2012.10.015

Cell

Transcription of Two Long Noncoding RNAs Mediates Mating-Type Control of Gametogenesis in Budding Yeast

Nature. 2012 Nov 15;491(7424):454-7. doi: 10.1038/nature11508. Epub 2012 Oct 14.

Long non-coding antisense RNA controls Uchl1 translation through an embedded SINEB2 repeat.

Epigenetic Regulation by Long Noncoding RNAs

Jeannie T. Lee

Science **338**, 1435 (2012);

DOI: 10.1126/science.1231776

Cell

A Long Noncoding RNA Controls Muscle Differentiation by Functioning as a Competing Endogenous RNA

Marcella Cesana,^{1,6} Davide Cacchiarelli,^{1,6} Ivano Legnini,¹ Tiziana Santini,¹ Olga Sthandier,¹ Mauro Chinappi,² Anna Tramontano,^{2,3,4} and Irene Bozzoni^{1,3,4,5,*}

The Emergence of lncRNAs in Cancer Biology

John R. Prensner and Arul M. Chinnaiyan

LETTER

doi:10.1038/nature17161

Biochim Biophys Acta. 2016 Jul;1863(7 Pt B):1953-60. doi: 10.1016/j.bbamcr.2015.11.026. Epub 2015 Nov 24.

Super-enhancer lncs to cardiovascular development and disease.

Our


Melanoma addiction to the long non-coding RNA SAMMSON

Roberto Vendramin^{1,2}, Marco Spinazzi², Patrick Laurette³, Mark Fiers², Jasper Wouters⁴, Enrico Radaelli⁵, Marina Leonelli^{8,9}, Katrien Vanderheyden^{8,9}, Aljosja Rogiers^{1,2}, Els Hermans¹⁰, Pieter Baatsen², Stein Aerts¹¹, Jan Van Aelst^{12,13}, Joost van den Oord⁴, Bart de Strooper², Irwin Davidson³, Denis L. J. Lafontaine¹⁴, Desompele^{8,9}, Pieter Mestdagh^{8,9*} & Jean-Christophe Marine^{1,2*}

Gene regulation by long non-coding RNAs and its biological functions


Luisa Statello^{1,2,6}, Chun-Jie Guo^{3,6}, Ling-Ling Chen^{3,4,5} and Maite Huarte^{1,2}

Reference annotation




- Gold standard for reference annotation in human and mouse
- **Manual curation** + evidence-based transcript models (RNASeq)

Genome annotation with short-read RNAseq



-  RNAs are fragmented
-  Assembly problem for both :
 - mRNAs and lncRNAs
 - novel isoforms of known genes and novel genes


Genome annotation with long-read RNAseq



-  "No need to reassemble reads"
-  Characterize and quantify "full length" transcripts (repeat-associated transcripts)
-  Direct exon connectivity
-  Epitranscriptomics: identification of RNA modifications (m6A)




Unfragmented view of the transcriptome



► LR-RNAseq is not totally long/full-length...

Long-read RNA-seq Genome Annotation Assessment Project



Challenge 1: transcript isoform detection with a high-quality genome
Challenge 2: transcript isoform quantification
Challenge 3: de novo transcript isoform identification

nature methods

Explore content ▾ About the journal ▾ Publish with us ▾

[nature](#) > [nature methods](#) > [registered report](#) > [article](#)

Registered Report | [Open access](#) | Published: 07 June 2024

Systematic assessment of long-read RNA-seq methods for transcript identification and quantification

[Francisco J. Pardo-Palacios](#), [Dingjie Wang](#), [Fairlie Reese](#), [Mark Diekhans](#), [Sílvia Carbonell-Sala](#), [Brian Williams](#), [Jane E. Loveland](#), [Maite De María](#), [Matthew S. Adams](#), [Gabriela Balderrama-Gutierrez](#), [Amit K. Behera](#), [Jose M. Gonzalez Martinez](#), [Toby Hunt](#), [Julien Lagarde](#), [Cindy E. Liang](#), [Haoran Li](#), [Marcus Jerryd Meade](#), [David A. Moraga Amador](#), [Andrey D. Prjibelski](#), [Inanc Birol](#), [Hamed Bostan](#), [Ashley M. Brooks](#), [Muhammed Hasan Çelik](#), [Ying Chen](#), ... [Angela N. Brooks](#) ✉ [+ Show authors](#)

[Nature Methods](#) **21**, 1349–1363 (2024) | [Cite this article](#)

49k Accesses | **101** Citations | **111** Altmetric | [Metrics](#)

● Long-read methods



StringTie

bambu




Bambu



IsoQuant

...


Long-read RNA-seq Genome Annotation Assessment Project



Challenge 1: transcript isoform detection with a high-quality genome
Challenge 2: transcript isoform quantification
Challenge 3: de novo transcript isoform identification



StringTie



StringTie



IsoQuant

...

nature methods

Explore content ▾ About the journal ▾ Publish with us ▾

[nature](#) > [nature methods](#) > [registered report](#) > [article](#)

Registered Report | [Open access](#) | Published: 07 June 2024

Systematic assessment of long-read RNA-seq methods for transcript identification and quantification

[Francisco J. Pardo-Palacios](#), [Dingjie Wang](#), [Fairlie Reese](#), [Mark Diekhans](#), [Sílvia Carbonell-Sala](#), [Brian Williams](#), [Jane E. Loveland](#), [Maite De María](#), [Matthew S. Adams](#), [Gabriela Balderrama-Gutierrez](#), [Amit K. Behera](#), [Jose M. Gonzalez Martinez](#), [Toby Hunt](#), [Julien Lagarde](#), [Cindy E. Liang](#), [Haoran Li](#), [Marcus Jerryd Meade](#), [David A. Moraga Amador](#), [Andrey D. Prjibelski](#), [Inanc Birol](#), [Hamed Bostan](#), [Ashley M. Brooks](#), [Muhammed Hasan Çelik](#), [Ying Chen](#), ... [Angela N. Brooks](#)

+ Show authors

[Nature Methods](#) **21**, 1349–1363 (2024) | [Cite this article](#)

49k Accesses | **101** Citations | **111** Altmetric | [Metrics](#)

❌ "Significant discrepancies among LR-RNAseq methods for transcript discovery."

➡ Better multiple tools

❌ No tool include lncRNA annotation (coding potential)

Long-read RNA sequencing : state of the art

**Multiple tools
for discovery**

**Annotate
lncRNAs**

**Transcript
classification**


**Full-length
assessment**

Long-read RNA sequencing : state of the art

	Multiple tools for discovery	Annotate lncRNAs	Transcript classification	Full-length assessment	Links
nf-core/nanoseq	 Bambu or Stringtie				https://nf-co.re/nanoseq/

nf-core/nanoseq

Nanopore demultiplexing, QC, alignment, and downstream analysis




Long-read RNA sequencing : state of the art


	Multiple tools for discovery	Annotate lncRNAs	Transcript classification	Full-length assessment	Links
nf-core/ nanoseq	 Bambu or Stringtie				https://nf-co.re/ nanoseq/
wf-human- transcriptome	 Stringtie only		 gffcompare		https://github.com/ epi2me-labs/wf- transcriptomes
TAGADA	 Stringtie only	 FEELnc	 FEELnc_classifier		https://github.com/ FAANG/analysis- TAGADA

Long-read RNA sequencing : state of the art


	Multiple tools for discovery	Annotate lncRNAs	Transcript classification	Full-length assessment	Links
nf-core/ nanoseq	 Bambu or Stringtie				https://nf-co.re/ nanoseq/
wf-human- transcriptome	 Stringtie only		 gffcompare		https://github.com/ epi2me-labs/wf- transcriptomes
TAGADA	 Stringtie only	 FEELnc	 FEELnc_classifier		https://github.com/ FAANG/analysis- TAGADA
ANNEXA					https://github.com/ IGDRion/ANNEXA/

Nicolaï
HOFFMANN






ANalysis with Nextflow for EXtended Annotation




IGDRion/ANNEXA

nextflow






ANalysis with Nextflow for EXtended Annotation



IGDRion/ANNEXA

nextflow






IGDRion/ANNEXA




- Bambu : better when reference annotation is more complete
- StringTie : better to identify novel isoforms




IGDRion/ANNEXA

- ▶ **FEELnc** to compute the coding potential of all novel transcripts from novel genes
 - ▶ **FEELnc** uses a RandomForest model trained with sequence features
 - ▶ ORF sizes
 - ▶ k-mer profiles
 - ▶ Transcript length
 - ▶ **TransDecoder** allows CDS, 5'UTR and 3'UTR in novel mRNAs
- ➡ Allows to discriminate between novel mRNAs versus lncRNAs




IGDRion/ANNEXA



- ➔ Novel isoform of known gene
- ➔ Extension of known gene
- ➔ antisense lncRNA
- ➔ Long intergenic ncRNA





<https://github.com/IGDRion/TransforKmers>




IGDRion/ANNEXA

- Transformer-based model to classify DNA/RNA sequences.
 - Pretrain with species-specific genomic sequences (DNABERT)
 - Finetuning with CAGE-validated TSSs vs Random sequences



TFK model






IGDRion/ANNEXA


► Automatically generated .pdf report at 3 levels genes/transcripts/exons

► Multiple features :


- Number of mRNAs versus lncRNAs
- Number of single-exon genes
- Breadth of expression /Tissue specificity
- Gene extensions



Proportion of mono versus multi-isoform genes



➡ Allows to QC novel annotated genes/transcripts/exons



IGDRion : <https://igdr.univ-rennes.fr/igdrion>



ENT EN FR Extranet Se connecter

igdr Institut de Génétique & Développement de Rennes
IGDR UMR 6290 CNRS - UR1


L'institut La recherche Orientations Scientifiques La formation **Services & équipements**

Accueil / Services & équipements / Séquençage

Séquençage de troisième génération

Séquençage longue lecture : Oxford Nanopore Technology (ONT)


La technologie de séquençage à lecture longue (Long-read sequencing) proposée par ONT (Oxford Nanopore Technologies) est basée sur la détection d'un courant différentiel lors du passage d'une molécule unique linéaire à travers un nanopore biologique ancré dans une membrane synthétique. Le nanopore est un capteur de flux ionique qui varie selon l'encombrement stérique du composé qui le traverse, ce qui permet d'identifier la séquence d'un brin d'ADN sur une grande longueur (jusqu'à plusieurs Mb) quelque soit sa composition en éléments répétés, en régions riche en GC ou méthylées.



Présentation de la technologie de séquençage troisième génération



Mise à Disposition (MAD)



Localisation



Formulaire de demande d'utilisation du séquenceur Nanopore



Contact & Equipe

Contacts



Thomas DERRIEN
Equipe Génétique du Chien
thomas.derrien@univ-rennes.fr
+33 (0)2 23 23 65 34



Edouard CADIEU
Equipe Génétique du Chien
edouard.cadieu@univ-rennes.fr
+33 (0)2 23 23 44 84

Ailleurs sur le web...

<https://nanoporetech.com/>

Voir aussi...

[Où sommes nous ?](#)



INPUT SAMPLES



LIBRARY PREP.




SEQUENCING





BIOINFORMATIC ANALYSES





➡ ~6M reads
➡ Variable number of reads per sample



➡ length~900nt
➡ Qscore ~10.5 (old chemistry)

Reference genome :



canFam4

Reference annotations (x3) :



Ensembl (Dyer et al., 2025)



RefSeq

Refseq (O'Leary et al., 2016)



Uppsala Univ (Wang et al., 2021)

Reference genome :



GRCh38

Reference annotations (x2):




Gencode (Mudge et al., 2025)



CHESS (Varabyou et al., 2023)




KNOWN



➡ Bambu reconstructs more **known** genes/transcripts than Stringtie consistently





KNOWN



- ➡ Bambu reconstructs more **known** genes/transcripts than Stringtie consistently
- ➡ Consistent pattern across the canine annotation



KNOWN





Annotation Type

Reconstructed
Reference (F)

- ➔ Bambu reconstructs more **known** genes/transcripts than Stringtie consistently
- ➔ Consistent pattern across the canine annotations
- ➔ As well as human annotations (although with lower rates)




NOVEL




- ➔ StringTie reconstructs more **novel** elements than Bambo
- ➔ Filtering process dramatically reduces number of novel elements



Bambu novel lncRNA



StringTie novel lncRNA



Different patterns in classification:



- **Bambu:** primarily **intergenic** (u) and **antisense** (x)
- **Stringtie:** **intergenic** (u) and **extension of known lncRNAs** (k), very few antisense



=> 5 novel genes
ortholog bw human and
dog (3 lncRNAs)


=> MM-specific

=> Potential candidates
for future functional
validation/biomarker



Conclusions

- Development of a new tool to process long-read RNA seq data and extend existing annotations in a **context-aware manner**.
- ANNEXA is **FAIR and freely available** (<https://github.com/IGDRion/ANNEXA/>)




- Applied ANNEXA on pilot comparative oncology long-read RNA seq, highlighting tool specific annotation of lncRNA biotypes



The cell as an RNA machine



Anastasia
RUSAKOVICH




What are circRNA?

- **circRNAs** are a type of non-coding RNA forming closed-loop structures


What are circRNA?

- **circRNAs** are a type of non-coding RNA forming closed-loop structures
- Highly **stable**, often **tissue-specific**, **potential disease biomarkers**
- Interest is growing due to their **roles in gene regulation and cancer**




How are they generated?

- Produced by **back-splicing**, a **non-canonical splicing event**
- **Multiple isoforms** can arise from the same linear RNA





How are they generated?

- Produced by **back-splicing**, a **non-canonical splicing event**
- **Multiple isoforms** can arise from the same linear RNA
- **Short-read sequencing struggles** to reconstruct full isoforms






Why long-read sequencing?

Oxford Nanopore allows sequencing of **full-length** molecules, capturing **internal structure** and **alternative splicing events**



Protocols for linearization of circRNAs





Fragmentation based approach



No repeats!



Rolling circle based approach




Have repeats!

Existing bioinformatic tools for circRNA long-read data

Step \ Tool	circNICK-lrs	isoCirc	CIRI-long
Expected Input	Fragmentation	Rolling circle amplification	Rolling circle reverse transcription
1 - Repeat Detection		Tandem Repeat Finder 	k-mer approach (k=8,11) 
2 - Mapping	PBLAT	minimap2	minimap2, SPOA, mappy, bwapy
3 - BSJ detection -Iterative alignment -Multi-tiered scoring -Spatial constraints -Splice site usage -Database reliance	✗ ✗ ✓ ✗ ✓ (3 databases)	✗ ✓ ✓ ✓ ✗ (only classif 2 db)	✓ ✗ ✗ ✓ ✓ (for BSJ correction, 1 db)

→ No comprehensive benchmark study comparing the bioinformatic tools developed for ONT sequencing data analysis

Our benchmark study



Requirements for simulation


Database features




circAtlas [Wu et al.], **circBase** [Glažar et al.]

An illustration of a person in a yellow shirt and blue pants holding a measuring tape.

**circRNA
lengths**



**circRNA type and
inner structure**




circRNA splice sites




Protocol features


Mouse brain, ONT (CIRI-long
protocol, SRA: CRR194180)




Rolling circles count and repeat pattern




Mature Length Distribution by circRNA Type





Exon Count Distribution by circRNA Type




Splice Site Distribution by circRNA Type






How many rolling
circles protocol
introduces?



1 repeat = 1 copy
of circRNA

Our benchmark study




**Oxford
Nanopore
reads
generation**


Using NanoSim tool
(Hafezqorani et al. Gigascience, 2020)

Benchmark dataset

NanoSim



Our benchmark study




**Tool
application
and evaluation**


**Descriptive and
performance
metrics**

Descriptive metrics: Number of circRNAs detected





**Total isoforms
detected:
2295/7503
(30.58%)**




Descriptive metrics: Length distribution



Descriptive metrics: circRNA type distribution







Descriptive metrics: Expression level (TPM)






Performance evaluation

Transcript
level
(not exon-aware)






Performance evaluation

Transcript level
(not exon-aware)



Block level
(exon-aware)



Recall/Sensitivity = $TP / (TP + FN)$


Precision $s = TP / (TP + FP)$

Successful tool prediction

Failed tool prediction


Performance Metrics Comparison: Exon-aware vs. Not exon-aware

Analysis based on simulated reads (overlap fraction = 0.75)




Performance Metrics: Tool Combinations

Analysis based on simulated reads (overlap fraction = 0.75)




CircRNA Analysis Tools: Time vs Memory Performance



Main Findings

- Each tool has distinct strengths: **isoCirc** (precision and resources), **circNick-Irs** (recall and length), **CIRI-long** (balance and type assignment)
- Only **11.85% overlap** between all three tools - highlighting complementary detection
- All tools show **bias toward highly expressed circRNAs** and **lower sensitivity** for **exact inner structure**

Availability



Simulation framework is available on git



Perspectives


 **Expand protocols** → tool assessment on different protocols and species

 **Containerized pipeline** → tools application and result combination

 **Real data application** → Melanoma & hepatocellular carcinoma data generated in IGDR and OSS

Acknowledgements

IGDRion




Edouard
CADIEU




Aurore
BESSON



Victor
LE BARS



Nicolai
HOFFMANN



Anastasia
RUSAKOVICH



> BIOLOGY
IN SILICO for
NON CODING
GENOME and
ONCOLOGY




Yuna
BLUM




Christophe
HELIGON

Canine Genetics Team @ IGDR



Catherine ANDRE

Genouest bioinformatic



JBISR is a one-day symposium dedicated to **bioinformatics**, organized as part of the BIS2 ([Biology In Silico](#)) transversal axis at IGDR. The program highlights **cutting-edge research** in **AI-driven analysis** and **multi-modal data integration** in **health** and **cancer**.

09:30 - 10:00 Welcoming participants

10:00 - 10:10 Introductory remarks

10:10 - 11:45 **Session 1 - Integration of Multi-Modal Data in Biology and Cancer**

10:10 - 11:00 Keynote Speaker 1



DR. ANDREA RAU

Université Paris-Saclay, INRAE, AgroParisTech

Title: Tackling the statistical challenges of multi-omic integration

Dr. Andrea Rau, a leading expert in biostatistics and computational biology, will discuss cutting-edge methodologies for multi-omics data integration and highlight applications in oncology.

11:00 - 11:45 **3 selected presentations** (15 minutes for each presentation)

11:45 - 12:45 *Lunch Break*

12:45 - 13:45 **Poster all Sessions** with Coffee

13:45 - 15:15 **Session 2 - AI for Sequence and Image Analyses**

13:45 - 14:25 Keynote Speaker 2



DR. JULIETTE GRIFFIÉ

Science for Life Laboratory, Stockholm University, Sweden

Title: Interpretable ML for data driven therapeutic design and diagnosis

Dr. Juliette Griffié specializes in computational imaging and AI-driven biomedical modeling. Her talk will cover recent advances in microscopy image analysis, generative AI, and explainable AI in immunology applications in oncology.

14:25 - 15:10 **3 selected presentations** (15 minutes for each presentation)

15:15 - 15:45 *Coffee Break*

15:45 - 17:15 **Session 3 - Applications of AI and integrative approaches in Personalized Medicine and Health**

15:45 - 16:25 Keynote Speaker 3



DR. RÉMY NICOLLE

Centre de Recherche sur l'Inflammation (INSERM, CNRS, Université Paris Cité) Paris

Title: Multimodal Tumor phenotyping for treatment recommendation in pancreatic cancer

Dr. Rémy Nicolle, a renowned expert in onco-genomics and pancreatic cancer research, will discuss emerging multi-modal integrative approaches and AI methodologies for personalized medicine, focusing on patient stratification, biomarker discovery, and personalized therapeutic strategies in pancreatic cancer.

16:25 - 17:15 **3 selected presentations** (15 minutes for each presentation)

17:15 - 17:30 **Closing Session - Future Directions in Bioinformatics**

Closing Remarks and Farewell

